

IOWA STATE UNIVERSITY

Digital Repository

Integrated Crop Management News

Agriculture and Natural Resources

4-17-2006

Alfalfa weevil--consider cutting as a management option

Marlin E. Rice

Iowa State University, merice@iastate.edu

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Entomology Commons](#)

Recommended Citation

Rice, Marlin E., "Alfalfa weevil--consider cutting as a management option" (2006). *Integrated Crop Management News*. Paper 1238.
<http://lib.dr.iastate.edu/cropnews/1238>

This Article is brought to you for free and open access by the Agriculture and Natural Resources at Digital Repository @ Iowa State University. It has been accepted for inclusion in Integrated Crop Management News by an authorized administrator of Digital Repository @ Iowa State University. For more information, please contact digirep@iastate.edu.

INTEGRATED CROP MANAGEMENT

Alfalfa weevil--consider cutting as a management option

Somewhere in Iowa this spring, alfalfa weevil populations will exceed economic thresholds and cause significant yield loss. Now is the time to be scouting alfalfa for this pest. I like to use a sweep net to simply determine if larvae have hatched in the field. Sweeping even a small patch of alfalfa with a net can reveal whether or not larvae are present. If no larvae are found, then move to the next field. However, if they are found in the net, the next step is the stem-sampling technique. This technique more accurately determines the population size and the potential for economic damage.

Collect 30 stems by holding the top of the plant with one hand and breaking the base of the stem with the other hand, or cut it with a knife. Holding the top of the stem prevents larvae from dropping from the stem when it is broken or cut at the base. Then place the stems (upside down) inside a white, 5-gallon bucket and beat them against the side. Large larvae are easily knocked loose and can be counted, but newly developing leaves must be pulled apart to find very small, newly hatched larvae hidden in the plant tip.



[1]

Cutting is a management option for alfalfa weevil larvae. (Marlin E. Rice)



©MARLIN E. RICE

[2]

Damage to alfalfa by weevil larvae “skeletonizes” the leaves with only the tougher leaf veins remaining. (Marlin E. Rice)

Alfalfa weevil larvae have a very dark head, almost black, and are pale green with a white stripe down the back. When the alfalfa weevil hatch, they are approximately 1/16 inch in length and may be light yellow. After feeding for several days, they turn green. They are 5/16 inch in length when fully grown.

Alfalfa weevil larvae may be confused with larvae of the clover leaf weevil, although these are much larger, have a light brown head, and often have the white stripe edged with pink. Clover leaf weevil larvae usually hide around the base of the plant during the day, feed mostly in lower leaves at night, and rarely cause economic yield losses. Clover leaf weevil larvae should not be counted as part of the alfalfa weevil sample.



[3]

Alfalfa weevil larva with black head and damage to alfalfa leaf. (Marlin E. Rice)



©MARLIN E. RICE

[4]

Clover leaf weevil larva with light brown head. (Marlin E. Rice)

Economic thresholds are shown in Table 1. Measure the plant height and then determine the average number of weevil larvae per stem, based upon a 30-stem count, before consulting Table 1. The economic threshold depends on crop height, estimated crop value, control costs, and the growing conditions stated in Table 1. If weevils exceed the economic threshold, then consider cutting the forage as the first management option. However, make certain that the hay is at the optimal developmental stage before cutting. If the hay is not ready to be cut in several days, then an insecticide might be necessary. Several insecticides labeled for weevils are listed in Table 2.

Table 1. Economic thresholds based on alfalfa weevil larvae per stem, calculated from a 30-stem sample*

**Plant
Height (Inches) \$40/Ton \$70/Ton \$100/Ton Management Decision**

4	1.8-2.8	0.8-1.3	0.6-0.8	Reevaluate in 4 days. If damage and larval numbers are increasing, a long-residual insecticide is recommended to prevent severe yield loss.
6	2.0-3.0	0.8-1.5	0.6-1.0	
8	2.2-3.2	0.9-1.7	0.7-1.2	
10	2.3-3.5	0.9-1.9	0.8-1.4	If alfalfa is in vegetative stages, a short-residual insecticide should be used.
12	2.4-3.8	1.0-2.2	0.9-1.6	
14	2.5-4.2	1.2-2.5	1.0-1.8	
16	2.6-4.6	1.5-2.8	1.1-2.0	If >60 percent of alfalfa is in the bud stage, harvest is recommended. Evaluate stubble after harvest. If not scheduled to be cut within 7-10 days, a short-residual insecticide is recommended.
18	2.7-5.0	1.7-3.1	1.2-2.3	
20	2.8-5.8	2.0-3.4	1.4-2.6	
>20	3.0-7.0	2.4-4.0	1.6-3.0	

*Use the smaller threshold if alfalfa is drought-stressed or control costs are relatively low (\$7-10 per acre). Use a larger threshold if rainfall is abundant, diseased larvae are present, or control costs are relatively high (\$11-14 per acre).

Table 2. Insecticides labeled for alfalfa weevil

Insecticide	Rate per Acre (High and Low Rates)	Harvest Interval (Days)
Baythroid 2 E	1.6-2.8 ounces	7
Furadan 4 F	0.5-2 pints	7-28
Lannate LV	3 pints	0
Lorsban 4 E	1-2 pints	14-21
Mustang Max	2.24-4.0 ounces	3
Pounce 3.2 EC	4-8 ounces	0-14
Sevin XLR+	3 pints	7
Warrior	2.56-3.84 ounces	7

This article originally appeared on pages 87-89 of the IC-496 (7) -- April 17, 2006 issue.

Source URL:

<http://www.ipm.iastate.edu/ipm/icm//ipm/icm/2006/4-17/alfalfa.html>

Links:

- [1] <http://www.ipm.iastate.edu/ipm/icm/node/395>
- [2] <http://www.ipm.iastate.edu/ipm/icm/node/394>
- [3] <http://www.ipm.iastate.edu/ipm/icm/node/66>
- [4] <http://www.ipm.iastate.edu/ipm/icm/node/397>

IOWA STATE UNIVERSITY
University Extension